

ON THE NUMBER OF POINTS OF THE LUSZTIG NILPOTENT VARIETY OVER FINITE FIELDS

O. SCHIFFMANN

ABSTRACT. We give a closed expression for the number of points over finite fields of the Lusztig nilpotent variety associated to any quiver without edge loops, in terms of Kac's A -polynomial. We conjecture a similar result for quivers in which edge loops are allowed. Finally, we give a formula for the number of points over a finite field of the various stratas of the Lusztig nilpotent variety involved in the geometric realization of the crystal graph.

1. STATEMENT OF THE RESULT

1.1. Let $Q = (I, H)$ be a finite quiver, with vertex set I and edge set H . For $h \in H$ we will denote by h', h'' the initial and terminal vertex of h . Note that we allow edge loops, i.e. edges h satisfying $h' = h''$. As usual we denote by

$$\langle \mathbf{d}, \mathbf{e} \rangle = \sum_{i \in I} d_i e_i - \sum_{h \in H} d_{h'} e_{h''}$$

the Euler form on $\mathbb{Z}^I$, and by $(\ , \)$ its symmetrized version : $(\mathbf{d}, \mathbf{e}) = \langle \mathbf{d}, \mathbf{e} \rangle + \langle \mathbf{e}, \mathbf{d} \rangle$. We also set $\mathbf{d} \cdot \mathbf{e} = \sum_i d_i e_i$. Let $\bar{Q} = (I, H \sqcup H^*)$ be the doubled quiver, obtained from Q by replacing each arrow h by a pair of arrows (h, h^*) going in opposite directions.

Fix a field k . For each dimension vector $\mathbf{d} \in \mathbb{N}^I$ we fix an I -graded k -vector space $V_{\mathbf{d}} = \bigoplus_i V_i$ and we set

$$E_{\mathbf{d}} = \bigoplus_{h \in H} \text{Hom}(V_{h'}, V_{h''}), \quad \bar{E}_{\mathbf{d}} = \bigoplus_{h \in H \sqcup H^*} \text{Hom}(V_{h'}, V_{h''}).$$

Elements of $E_{\mathbf{d}}$ (resp. $\bar{E}_{\mathbf{d}}$) will be denoted by $(x_h)_h$ (resp. by $(x_h, x_{h^*})_h$). We say that $(x_h)_h$ (resp. $(x_h, x_{h^*})_h$) is nilpotent if there exists an I -graded flag

$$L^\bullet = (\{0\} = L^0 \subset L^1 \subset \dots \subset L^{|\mathbf{d}|} = V_{\mathbf{d}}), \quad |\mathbf{d}| = \sum_i d_i$$

of vector subspaces in $V_{\mathbf{d}}$ such that

$$x_h(L^l) \subseteq L^{l-1}, \quad l = 0, \dots, |\mathbf{d}|, \quad h \in H$$

(resp. such that

$$x_h(L^l) \subseteq L^{l-1}, \quad x_{h^*}(L^l) \subseteq L^l, \quad l = 0, \dots, |\mathbf{d}|, \quad h \in H).$$

The sets of nilpotent representations form Zariski closed subvarieties $E_{\mathbf{d}}^{\text{nil}} \subseteq E_{\mathbf{d}}$ and $\bar{E}_{\mathbf{d}}^{\text{nil}} \subseteq \bar{E}_{\mathbf{d}}$. When Q has no edge loops, the above nilpotency condition for elements in $\bar{E}_{\mathbf{d}}$ is equivalent to the standard nilpotency condition (i.e. that there exists an integer $N > 0$ such that the composition $x_{h_1} x_{h_2} \dots x_{h_N}$ vanishes for any collection of edges $h_1, \dots, h_N$ in $H \sqcup H^*$).

The group $G_{\mathbf{d}} = \prod_i GL(V_i)$ acts on $E_{\mathbf{d}}$ and $\bar{E}_{\mathbf{d}}$ by conjugation, and preserves the subvarieties $E_{\mathbf{d}}^{\text{nil}}, \bar{E}_{\mathbf{d}}^{\text{nil}}$.

Let us now assume that $k = \mathbb{C}$. The trace map

$$Tr : \overline{E}_{\mathbf{d}} \rightarrow \mathbb{C}, \quad (x_h, x_{h^*})_h \mapsto \sum_h tr(x_h x_{h^*})$$

identifies $\overline{E}_{\mathbf{d}}$ with $T^*E_{\mathbf{d}}$. We set $\mathfrak{g}_{\mathbf{d}} = Lie(G_{\mathbf{d}}) = \bigoplus_i \mathfrak{gl}(V_i)$ and identify $\mathfrak{g}_{\mathbf{d}}$ with its dual $\mathfrak{g}_{\mathbf{d}}^*$ via the standard Killing form. The moment map for the action of $G_{\mathbf{d}}$ on $T^*E_{\mathbf{d}}$ may be written as

$$\mu_{\mathbf{d}} : \overline{E}_{\mathbf{d}} \rightarrow \mathfrak{g}_{\mathbf{d}}, \quad (x_h, x_{h^*})_h \mapsto \sum_{h \in H} [x_h, x_{h^*}].$$

Following Lusztig (see [L1]) we consider the intersection

$$\Lambda_{\mathbf{d}} = \mu_{\mathbf{d}}^{-1}(0) \cap \overline{E}_{\mathbf{d}}^{nil} \subset \overline{E}_{\mathbf{d}}.$$

This variety is often called the *Lusztig nilpotent variety* (see [B] when Q has edge loops). It is a closed subvariety of $\overline{E}_{\mathbf{d}}$, which, in general, possesses many irreducible components and is singular. In addition, $\Lambda_{\mathbf{d}}$ is a Lagrangian subvariety of $\overline{E}_{\mathbf{d}}$ (see [L1], [B]). It plays an important role in the geometric approach to quantum groups and crystal graphs based on quiver varieties (see [KS] or [S, Lect. 4]).

1.2. The map μ and hence the variety $\Lambda_{\mathbf{d}}$ are defined over an arbitrary field k . The aim of this note is to establish a formula for the number of points of $\Lambda_{\mathbf{d}}$ over finite fields, in terms of Kac's A -polynomial ([K], see below). We will give a formula for the generating series of the $|\Lambda_{\mathbf{d}}(\mathbb{F}_q)|$. Before we can state our result, we need to fix a few notations. Let us consider the space

$$\mathbf{L} = \mathbb{Q}(t)[[z_i]]_{i \in I}$$

of power series in variables $z_i, i \in I$, with coefficients in the field $\mathbb{Q}(t)$. Here t is a formal variable. For $\mathbf{d} \in \mathbb{N}^I$ we write $z^{\mathbf{d}} = \prod_i z_i^{d_i}$. Let $Exp : \mathbf{L} \rightarrow \mathbf{L}$ be the λ -ring version of the exponential map, i.e. $Exp(f) = exp(\sum_k \frac{1}{k} \psi_k(f))$, where $\psi_k : \mathbf{L} \rightarrow \mathbf{L}$ is the k th Adams operator satisfying $\psi_k(z^{\mathbf{d}}) = z^{k\mathbf{d}}, \psi_k(t^l) = t^{kl}$. For instance, we have $Exp(z^{\mathbf{d}}) = 1/(1 - z^{\mathbf{d}})$.

1.3. Let $A_{\mathbf{d}}(t)$ be Kac's A -polynomial attached to the quiver Q and to the dimension vector $\mathbf{d}$, i.e. for any finite field $\mathbb{F}_q$, the integer $A_{\mathbf{d}}(q)$ is equal to the number of isomorphism classes of absolutely indecomposable representation of Q of dimension $\mathbf{d}$ over $\mathbb{F}_q$. The existence of $A_{\mathbf{d}}(t)$ is due to Kac ([K]), as is the fact that $A_{\mathbf{d}}(t) \in \mathbb{Z}[t]$ is unitary of degree $1 - \langle \mathbf{d}, \mathbf{d} \rangle$. The fact that $A_{\mathbf{d}}(t)$ has positive coefficients was only recently proved in [HLR-V]. By Kac's theorem, $A_{\mathbf{d}}(t) = 0$ unless $\mathbf{d}$ belongs to the set Δ^+ of positive roots of Q . Let us denote by $\overline{A}_{\mathbf{d}}(t) = A_{\mathbf{d}}(t^{-1})t^{1 - \langle \mathbf{d}, \mathbf{d} \rangle}$ the reciprocal polynomial of $A_{\mathbf{d}}(t)$. Let us consider the element

$$(1.1) \quad P_Q(t, z) := Exp\left(\frac{1}{t-1} \sum_{\mathbf{d}} \overline{A}_{\mathbf{d}}(t) t^{\langle \mathbf{d}, \mathbf{d} \rangle} z^{\mathbf{d}}\right) = Exp\left(\frac{1}{1-t^{-1}} \sum_{\mathbf{d}} A_{\mathbf{d}}(t^{-1}) z^{\mathbf{d}}\right) \in \mathbf{L}.$$

If $A_{\mathbf{d}}(t) = \sum_n a_{\mathbf{d},n} t^n$, the definition of $P_Q(t)$ may be rewritten as follows :

$$P_Q(t, z) = \prod_{\mathbf{d} \in \Delta^+} \prod_{l \geq 0} \prod_{n = \langle \mathbf{d}, \mathbf{d} \rangle - 1}^0 \frac{1}{(1 - t^{n-l} z^{\mathbf{d}})^{a_{\mathbf{d}, -n}}}.$$

Observe that the Fourier modes of $P_Q(t, z)$ are all rational functions in t regular outside of $t = 1$ and this allows us to evaluate $P_Q(t, z)$ at any $t \neq 1$.

We may now state our main (and only) result :

Theorem 1.1. *Assume that Q has no edge loops and let us set*

$$\lambda_Q(q, z) = \sum_{\mathbf{d}} \frac{|\Lambda_{\mathbf{d}}(\mathbb{F}_q)|}{|G_{\mathbf{d}}(\mathbb{F}_q)|} q^{\langle \mathbf{d}, \mathbf{d} \rangle} z^{\mathbf{d}} \in \mathbb{Q}[[z_i]]_i.$$

Then

$$(1.2) \quad \lambda_Q(q, z) = P_Q(q, z).$$

Conjecture 1.2. *Theorem 1.1 holds without any assumption on Q .*

Remarks. i) Set $\zeta_{\mathbf{d}}(q, u) = \exp\left(\sum_{l \geq 1} A_{\mathbf{d}}(q^l) \frac{u^l}{l}\right)$. This may be thought of as the 'zeta function' of the quotient

$$\mathcal{M}_{\mathbf{d}}^{abs.ind} = \{(x_h)_h \in E_{\mathbf{d}} \mid (x_h \otimes \overline{\mathbb{F}_q})_h \text{ indecomposable}\} / G_{\mathbf{d}}.$$

Of course, $\mathcal{M}_{\mathbf{d}}^{abs.ind}$ is not an algebraic variety in any sense – it is only a constructible subset of the stack $\mathcal{M}_{\mathbf{d}} = E_{\mathbf{d}} / G_{\mathbf{d}}$ classifying representations of Q of dimension $\mathbf{d}$, and we can not speak of its zeta function in any precise sense. Using this notation, we may restate our result as the equality

$$\lambda_Q(q, z) = \prod_{\mathbf{d} \in \Delta^+} \prod_{l \geq 0} \zeta_{\mathbf{d}}(q^{-1}, q^{-l} z^{\mathbf{d}}).$$

ii) It is possible to give a closed expression for the power series $P_Q(t, z)$, as an immediate consequence of Hua's formula, see (2.5).

1.4. Let us provide a few simple examples for Theorem 1.1 and Conjecture 1.2.

i) Assume that Q is a finite Dynkin quiver. Then $A_{\mathbf{d}} = 1$ for all $\mathbf{d} \in \Delta^+$, and thus

$$\lambda_Q(q, z) = \prod_{\mathbf{d} \in \Delta^+} \prod_{l \geq 0} \frac{1}{1 - q^{-l} z^{\mathbf{d}}} = \exp\left(\frac{q}{q-1} \sum_{\mathbf{d} \in \Delta^+} z^{\mathbf{d}}\right).$$

ii) Now let us assume that Q is an affine quiver. Then $\Delta^+ = \Delta_{im}^+ \sqcup \Delta_{re}^+$ with

$$\Delta_{im}^+ = \mathbb{N}_{\geq 1} \delta, \quad \Delta_{re}^+ = \{\Delta_0^+ + \mathbb{N} \delta\} \sqcup \{\Delta_0^- + \mathbb{N}_{\geq 1} \delta\}$$

where δ is the minimal imaginary root, and where Δ_0 is the root system of an underlying finite type subquiver $Q_0 \subset Q$. We have $A_{\mathbf{d}}(t) = 1$ for $\mathbf{d} \in \Delta_{re}^+$ while $A_{\mathbf{d}}(T) = t + r$, with $r = \text{rank}(Q_0) = |I| - 1$. This yields

$$\lambda_Q(q, z) = \exp\left(\frac{\sum_{\mathbf{d} \in \Delta_0^+} q z^{\mathbf{d}} + (1 + rq) z^{\delta} + \sum_{\mathbf{d} \in \Delta_0^-} q z^{\mathbf{d} + \delta}}{(q-1)(1-z^{\delta})}\right).$$

iii) Finally, consider the Jordan quiver Q with one vertex and one loop. Then $\Delta^+ = \mathbb{N}_{\geq 1}$ and $A_{\mathbf{d}}(t) = t$ for all $\mathbf{d} \geq 1$. In this case, $\Lambda_{\mathbf{d}}$ is just the variety $\text{Com}_{\mathfrak{gl}_{\mathbf{d}}}^{\bullet, nil}$ of pairs of commuting $\mathbf{d} \times \mathbf{d}$ -matrices, with the second matrix being nilpotent. Theorem 1.1 is not applicable here, but a direct computation (see [R-V]) yields, in accordance with Conjecture 1.2

$$\sum_{\mathbf{d} \geq 0} \frac{|\text{Com}_{\mathfrak{gl}_{\mathbf{d}}}^{\bullet, nil}(\mathbb{F}_q)|}{|GL_{\mathbf{d}}(\mathbb{F}_q)|} z^{\mathbf{d}} = \prod_{\mathbf{d} \geq 1} \prod_{l \geq 0} \frac{1}{1 - q^{-l} z^{\mathbf{d}}} = \exp\left(\frac{qz}{(q-1)(1-z)}\right).$$

The above formula should be compared to the Feit-Thompson formula for the number of points in the (non-nilpotent) commuting varieties over finite fields (see [FT]).

Remark. The same techniques allow one to prove the following similar formula

$$\sum_{\mathbf{d} \geq 0} \frac{|\text{Com}_{\mathfrak{gl}_{\mathbf{d}}}^{nil, nil}(\mathbb{F}_q)|}{|GL_{\mathbf{d}}(\mathbb{F}_q)|} z^{\mathbf{d}} = \prod_{\mathbf{d} \geq 1} \prod_{l \geq 1} \frac{1}{1 - q^{-l} z^{\mathbf{d}}} = \exp\left(\frac{z}{(q-1)(1-z)}\right).$$

where $\text{Com}_{\mathfrak{gl}_{\mathbf{d}}}^{nil, nil}$ is now the variety of pairs of commuting nilpotent matrices. One might ask if there is an analogous formula for an arbitrary quiver (allowing edge loops), when one counts

the number of points of the nilpotent variety $\Lambda_{\mathbf{d}}^{nil} \subset \Lambda_{\mathbf{d}}$ defined using the *standard* nilpotency condition. In view of Conjecture 1.2, one might expect that

$$\sum_{\mathbf{d}} \frac{|\Lambda_{\mathbf{d}}(\mathbb{F}_q)|}{|G_{\mathbf{d}}(\mathbb{F}_q)|} q^{\langle \mathbf{d}, \mathbf{d} \rangle} z^{\mathbf{d}} = \text{Exp} \left(\frac{1}{t-1} \sum_{\mathbf{d}} \bar{A}_{\mathbf{d}}^{nil}(t) t^{\langle \mathbf{d}, \mathbf{d} \rangle} z^{\mathbf{d}} \right)$$

where $\bar{A}_{\mathbf{d}}^{nil}(t)$ counts the number of absolutely indecomposable representations of Q of dimension $\mathbf{d}$ in which all edge loops act nilpotently.

1.5. We finish the section with two short remarks.

Poincaré duality. It is not difficult to show that

$$(1.3) \quad \sum_{\mathbf{d}} \frac{|\mu_{\mathbf{d}}^{-1}(0)(\mathbb{F}_q)|}{|G_{\mathbf{d}}(\mathbb{F}_q)|} q^{\langle \mathbf{d}, \mathbf{d} \rangle} z^{\mathbf{d}} = \text{Exp} \left(\frac{q}{q-1} \sum_{\mathbf{d}} A_{\mathbf{d}}(q) z^{\mathbf{d}} \right)$$

(see [M2, Thm.5.1] for a similar result in the context of Donaldson-Thomas theory). The stack quotient $[\Lambda_{\mathbf{d}}/G_{\mathbf{d}}]$ is Lagrangian inside $[\mu_{\mathbf{d}}^{-1}(0)/G_{\mathbf{d}}]$. Comparing (1.3) with (1.1) we see that, as far as point counting goes, $[\Lambda_{\mathbf{d}}/G_{\mathbf{d}}]$ and $[\mu_{\mathbf{d}}^{-1}(0)/G_{\mathbf{d}}]$ are in some sense Poincaré dual to each other. In fact, our proof of Theorem 1.1 uses such a Poincaré duality for appropriate *stable* (in particular smooth) quotients of $\Lambda_{\mathbf{d}}$ and $\mu_{\mathbf{d}}^{-1}(0)$, see Section 2.1. It would be interesting to be able to deduce the relation between the point counts of $\Lambda_{\mathbf{d}}$ and $\mu_{\mathbf{d}}^{-1}(0)$ directly from geometric properties of the stacks $[\Lambda_{\mathbf{d}}/G_{\mathbf{d}}]$ and $[\mu_{\mathbf{d}}^{-1}(0)/G_{\mathbf{d}}]$.

Kac's conjecture. Let Q be a quiver without edge loops and let $\mathfrak{g}$ be the associated Kac-Moody algebra. By a theorem of Kashiwara-Saito ([KS], conjectured by Lusztig in [L2]), the number of irreducible components of $\Lambda_{\mathbf{d}}$ is the dimension of the $\mathbf{d}$ root space in the envelopping algebra $U^+(\mathfrak{g})$. By the Lang-Weil theorem, we have

$$|\Lambda_{\mathbf{d}}(\mathbb{F}_q)| = |\text{Irr}(\Lambda_{\mathbf{d}})| q^{\dim(\Lambda_{\mathbf{d}})} + O(q^{\dim(\Lambda_{\mathbf{d}})-1/2})$$

from which it follows that

$$\frac{|\Lambda_{\mathbf{d}}(\mathbb{F}_q)|}{|G_{\mathbf{d}}(\mathbb{F}_q)|} q^{\langle \mathbf{d}, \mathbf{d} \rangle} = |\text{Irr}(\Lambda_{\mathbf{d}})| + O(q^{-1/2}) = \dim(U^+(\mathfrak{g})[\mathbf{d}]) + O(q^{-1/2})$$

which we may write as

$$(1.4) \quad \lambda_Q(q, z) = \sum_{\mathbf{d}} \dim(U^+(\mathfrak{g})[\mathbf{d}]) z^{\mathbf{d}} + O(q^{-1/2}).$$

On the other hand, by Theorem 1.1 we have

$$(1.5) \quad \lambda_Q(q, z) = \text{Exp} \left(\sum_{\mathbf{d}} a_{\mathbf{d},0} z^{\mathbf{d}} \right) + O(q^{-1/2}) = \prod_{\mathbf{d}} \frac{1}{(1 - z^{\mathbf{d}})^{a_{\mathbf{d},0}}} + O(q^{-1/2}).$$

By the PBW theorem, $\sum_{\mathbf{d}} \dim(U^+(\mathfrak{g})[\mathbf{d}]) z^{\mathbf{d}} = \prod_{\mathbf{d}} (1 - z^{\mathbf{d}})^{-\dim(\mathfrak{g}[\mathbf{d}])}$. Combining (1.4) and (1.5) we obtain that $a_{\mathbf{d},0} = \dim(\mathfrak{g}[\mathbf{d}])$, which is the statement of Kac's conjecture. This conjecture was proved by Hausel in [Ha], using his computation of the Betti numbers of Nakajima quiver varieties. Note that our derivation of Theorem 1.1 uses Hausel's result in a crucial manner, so that the above is not a new proof of Kac's conjecture but rather a reformulation of Hausel's proof in terms of Lusztig nilpotent varieties instead of Lagrangian Nakajima quiver varieties (more in the spirit of [CBVdB]).

2. PROOF OF THE THEOREM

2.1. We assume henceforth that the quiver Q has no edge loops. Let $\mathbf{k}$ be a field. We recall the definition of Nakajima quiver varieties and state some of their properties (see [N1] for details). Fix $\mathbf{v}, \mathbf{w} \in \mathbb{N}^I$, let V, W be I -graded $\mathbf{k}$ -vector spaces of respective dimensions $\mathbf{v}, \mathbf{w}$ and set

$$M(\mathbf{v}, \mathbf{w}) = \bigoplus_{h \in H \sqcup H^*} \text{Hom}(V_{h'}, V_{h''}) \oplus \bigoplus_{i \in I} \text{Hom}(V_i, W_i) \oplus \bigoplus_{i \in I} \text{Hom}(W_i, V_i).$$

Elements of $M(\mathbf{v}, \mathbf{w})$ will be denoted $(\underline{x}, \underline{p}, \underline{q})$. The space $M(V, W)$ carries a natural symplectic structure, and the group $G_{\mathbf{v}} = \prod_i GL(V_i)$ acts in a Hamiltonian fashion. The associated moment map may be written as follows :

$$(2.1) \quad \begin{aligned} \mu : M(\mathbf{v}, \mathbf{w}) &\longrightarrow \bigoplus_i \mathfrak{gl}(V_i) \\ (\underline{x}, \underline{p}, \underline{q}) &\mapsto \left(\sum_{\substack{h \in H \\ h' = i}} x_{h^*} x_h - \sum_{\substack{h \in H \\ h'' = i}} x_h x_{h^*} + q_i p_i \right)_i \end{aligned}$$

The (categorical) symplectic quotient of $M(\mathbf{v}, \mathbf{w})$ by $G_{\mathbf{v}}$ is the variety $\mathcal{M}_0(\mathbf{v}, \mathbf{w}) = \mu^{-1}(0) // G_{\mathbf{v}}$. It is an affine variety which is singular in general. We say that an element $(\underline{x}, \underline{p}, \underline{q}) \in \mu^{-1}(0)$ is semistable if the following condition is satisfied :

$$\left(V' \subset \bigoplus_i \text{Ker}(q_i) \text{ and } \underline{x}(V') \subset V' \right) \Rightarrow V' = \{0\}.$$

We denote by $\mu^{-1}(0)^s$ the open subset of $\mu^{-1}(0)$ consisting of semistable points. The (geometric) quotient $\mathcal{M}(\mathbf{v}, \mathbf{w}) = \mu^{-1}(0)^s / G_{\mathbf{v}}$ is a smooth symplectic quasiprojective variety, and there is a natural projective morphism $\pi : \mathcal{M}(\mathbf{v}, \mathbf{w}) \rightarrow \mathcal{M}_0(\mathbf{v}, \mathbf{w})$. The dimension of $\mathcal{M}(\mathbf{v}, \mathbf{w})$ is equal to $2d(\mathbf{v}, \mathbf{w}) = 2\mathbf{v} \cdot \mathbf{w} - (\mathbf{v}, \mathbf{v})$. We put $\mathcal{L}(\mathbf{v}, \mathbf{w}) = \pi^{-1}(0)$. It is known that

$$\mathcal{L}(\mathbf{v}, \mathbf{w}) = \{G_{\mathbf{v}} \cdot (\underline{x}, \underline{p}, \underline{q}) \in \mathcal{M}(\mathbf{v}, \mathbf{w}) \mid \underline{p} = 0, \underline{x} \text{ is nilpotent}\}$$

and that $\mathcal{L}(\mathbf{v}, \mathbf{w})$ is Lagrangian in $\mathcal{M}(\mathbf{v}, \mathbf{w})$. When we want to specify the field over which we consider $\mathcal{M}(\mathbf{v}, \mathbf{w})$ or $\mathcal{L}(\mathbf{v}, \mathbf{w})$ we write $\mathcal{M}(\mathbf{v}, \mathbf{w})/\mathbf{k}, \mathcal{L}(\mathbf{v}, \mathbf{w})/\mathbf{k}$. In fact, the definitions of $\mathcal{M}(\mathbf{v}, \mathbf{w})$ and $\mathcal{L}(\mathbf{v}, \mathbf{w})$ make sense over any commutative ring R and we will use the notation $\mathcal{M}(\mathbf{v}, \mathbf{w})/R, \mathcal{L}(\mathbf{v}, \mathbf{w})/R$ for the corresponding R -schemes.

2.2. We assume that $\mathbf{k} = \mathbb{C}$ in this paragraph. The varieties $\mathcal{L}(\mathbf{v}, \mathbf{w})/\mathbb{C}$ and $\mathcal{M}(\mathbf{v}, \mathbf{w})/\mathbb{C}$ are homotopic. The Betti numbers of $\mathcal{M}(\mathbf{v}, \mathbf{w})/\mathbb{C}$ have been computed by T. Hausel (see [Ha], and see [M1] for another proof). Before we state this result we need a few notations. For $n \in \mathbb{N}$ and $\mathbf{n} = (n_i)_i \in \mathbb{N}^I$ we set

$$[\infty, n] = \prod_{k=1}^n (1 - t^k)^{-1}, \quad [\infty, \mathbf{n}] = \prod_i [\infty, n_i].$$

Let $\boldsymbol{\tau} = (\tau^i)_i$ be a set of partitions indexed by I . We put

$$|\boldsymbol{\tau}| = \sum_i |\tau^i|, \quad \boldsymbol{\tau}_l = (\tau_l^i)_i \in \mathbb{N}^I$$

and

$$X(\boldsymbol{\tau}, t) = \prod_k t^{\langle \boldsymbol{\tau}_k, \boldsymbol{\tau}_k \rangle} [\infty, \boldsymbol{\tau}_k - \boldsymbol{\tau}_{k+1}].$$

Define a power series in $\mathbf{L}$ depending on a vector $\mathbf{w} \in \mathbb{Z}^I$ as

$$r(\mathbf{w}, t, z) = \sum_{\boldsymbol{\tau}} t^{\mathbf{w} \cdot \boldsymbol{\tau}_1} X(\boldsymbol{\tau}, t^{-1}) z^{|\boldsymbol{\tau}|}.$$

Theorem 2.1 (Hausel). *The varieties $\mathcal{M}(\mathbf{v}, \mathbf{w})/\mathbb{C}$ have a pure Hodge structure, vanishing odd cohomology, and their Poincaré polynomials for compactly supported cohomology*

$$P_c(\mathcal{M}(\mathbf{v}, \mathbf{w})/\mathbb{C}, t) := \sum_i \dim H_c^{2i}(\mathcal{M}(\mathbf{v}, \mathbf{w})/\mathbb{C}, \mathbb{C}) t^i$$

are determined by the following formula :

$$(2.2) \quad \sum_{\mathbf{v}} t^{-d(\mathbf{v}, \mathbf{w})} P_c(\mathcal{M}(\mathbf{v}, \mathbf{w})/\mathbb{C}, t) z^{\mathbf{v}} = \frac{r(\mathbf{w}, t, z)}{r(0, t, z)}.$$

Proposition 2.2. *The varieties $\mathcal{L}(\mathbf{v}, \mathbf{w})/\mathbb{C}$ have a vanishing odd cohomology, and their Poincaré polynomials are determined by the following formula :*

$$(2.3) \quad \sum_{\mathbf{v}} t^{-d(\mathbf{v}, \mathbf{w})} P_c(\mathcal{L}(\mathbf{v}, \mathbf{w})/\mathbb{C}, t) z^{\mathbf{v}} = \frac{r(\mathbf{v}, t^{-1}, z)}{r(0, t^{-1}, z)}.$$

Proof. By Poincaré duality, we have

$$\begin{aligned} \dim H_c^{2i}(\mathcal{M}(\mathbf{v}, \mathbf{w})/\mathbb{C}, \mathbb{C}) &= \dim H^{4d(\mathbf{v}, \mathbf{w})-2i}(\mathcal{M}(\mathbf{v}, \mathbf{w})/\mathbb{C}, \mathbb{C}) \\ &= \dim H^{4d(\mathbf{v}, \mathbf{w})-2i}(\mathcal{L}(\mathbf{v}, \mathbf{w})/\mathbb{C}, \mathbb{C}) \\ &= \dim H_c^{4d(\mathbf{v}, \mathbf{w})-2i}(\mathcal{L}(\mathbf{v}, \mathbf{w})/\mathbb{C}, \mathbb{C}). \end{aligned}$$

The second and third equalities respectively come from the facts that $\mathcal{M}(\mathbf{v}, \mathbf{w})/\mathbb{C}$ and $\mathcal{L}(\mathbf{v}, \mathbf{w})/\mathbb{C}$ are homotopic and that the latter is projective. We deduce that

$$(2.4) \quad P_c(\mathcal{L}(\mathbf{v}, \mathbf{w})/\mathbb{C}, t) = t^{2d(\mathbf{v}, \mathbf{w})} P_c(\mathcal{M}(\mathbf{v}, \mathbf{w})/\mathbb{C}, t^{-1}).$$

Relation (2.3) is thus a consequence of (2.2). $\square$

The relation between the function $r(\mathbf{w}, t, z)$ and the collection of Kac polynomials is given by the following result of Hua, see [Hu] :

Theorem 2.3 (Hua). *We have*

$$(2.5) \quad r(0, t, z) = \text{Exp} \left(\frac{1}{t-1} \sum_{\mathbf{d}} A_{\mathbf{d}}(t) z^{\mathbf{d}} \right)$$

2.3. We now consider the quiver varieties $\mathcal{L}(\mathbf{v}, \mathbf{w})/\mathbb{F}_q$ and $\mathcal{M}(\mathbf{v}, \mathbf{w})/\mathbb{F}_q$ over finite fields. We say that a variety X defined over a finite field $\mathbb{F}_q$ is strictly polynomial count if there exists a polynomial $P(t) \in \mathbb{Q}[t]$ such that for any $r \geq 1$ we have $|X(\mathbb{F}_{q^r})| = P(q^r)$.

Proposition 2.4. *The variety $\mathcal{L}(\mathbf{v}, \mathbf{w})/\mathbb{F}_q$ is pure and strictly polynomial count over any finite field $\mathbb{F}_q$. It has no odd (étale) cohomology. Its counting polynomial is equal to the Poincaré polynomial $P_c(\mathcal{L}(\mathbf{v}, \mathbf{w})/\mathbb{C}, t)$.*

Proof. A similar statement is proved by Nakajima in [N3, Sec. 5–8] for virtual Hodge polynomials, for ADE quivers. Our method is an adaptation of his.

Consider the following $\mathbb{G}_m$ -action on $\mathcal{M}(\mathbf{v}, \mathbf{w})$:

$$t \cdot (\underline{x}, \underline{p}, \underline{q}) = (t\underline{x}, t\underline{p}, t\underline{q}).$$

There is a compatible action of $\mathbb{G}_m$ on the affine quotient $\mathcal{M}_0(\mathbf{v}, \mathbf{w})$.

Lemma 2.5. *Let R be any commutative ring. The above $\mathbb{G}_m$ -action induces Bialynicki-Birula type decompositions*

$$(2.6) \quad \mathcal{M}(\mathbf{v}, \mathbf{w})/R = \bigsqcup_{\rho} U_{\rho}/R, \quad \mathcal{L}(\mathbf{v}, \mathbf{w})/R = \bigsqcup_{\rho} V_{\rho}/R$$

where U_{ρ}/R and V_{ρ}/R are affine fiber bundles over a smooth projective variety $\mathcal{M}_{\rho}/R$. In addition, we have

$$(2.7) \quad \mathcal{M}_{\rho}/\mathbb{C} \simeq (\mathcal{M}_{\rho}/\mathbb{Z}) \otimes \mathbb{C}, \quad \mathcal{M}_{\rho}/\mathbb{F}_q \simeq (\mathcal{M}_{\rho}/\mathbb{Z}) \otimes \mathbb{F}_q.$$

Proof. Since the $\mathbb{G}_m$ -action on $\mathcal{M}_0(\mathbf{v}, \mathbf{w})/R$ contracts everything to 0 and since $\pi^{-1}(0) = \mathcal{L}(\mathbf{v}, \mathbf{w})/R$ is projective the $\mathbb{G}_m$ -action on $\mathcal{M}(\mathbf{v}, \mathbf{w})$ is contracting as well. From the definition of $\mathcal{M}(\mathbf{v}, \mathbf{w})$ as a G.I.T quotient (see e.g. [N1, (3.6)]) we see that $\mathcal{M}(\mathbf{v}, \mathbf{w})/R$, and hence also $\mathcal{L}(\mathbf{v}, \mathbf{w})/R$ may be covered by $\mathbb{G}_m$ -invariant affine subschemes. We may thus apply Hesselink's version of the Bialynicki-Birula theorem [He, Thm. 5.8] to deduce that the fixed point scheme $\mathcal{M}(\mathbf{v}, \mathbf{w})^{\mathbb{G}_m}/R$ is regular and that there is a decomposition $\mathcal{M}(\mathbf{v}, \mathbf{w})/R = \bigsqcup_{\rho} U_{\rho}/R$ as in (2.6). Notice that $\mathcal{M}(\mathbf{v}, \mathbf{w})^{\mathbb{G}_m}/R \subset \mathcal{L}(\mathbf{v}, \mathbf{w})/R$ hence each connected component $\mathcal{L}_{\rho}/R$ of $\mathcal{M}(\mathbf{v}, \mathbf{w})^{\mathbb{G}_m}/R$ is projective. Applying Hesselink's theorem to the inverse $\mathbb{G}_m$ -action $t \star (\underline{x}, \underline{p}, \underline{q}) = t^{-1} \cdot (\underline{x}, \underline{p}, \underline{q})$ yields the decomposition in (2.6) for $\mathcal{L}(\mathbf{v}, \mathbf{w})/R$. Observe that the $\star$ -action of $\mathbb{G}_m$ on $\mathcal{M}_0(\mathbf{v}, \mathbf{w})/R$ is dilating hence the attracting scheme (called concentrator scheme in [He]) is equal to $\pi^{-1}(0) = \mathcal{L}(\mathbf{v}, \mathbf{w})/R$. The assertions in (2.7) come from the fact that taking $\mathbb{G}_m$ -invariants commutes with base change. $\square$

We may now finish the proof of Proposition 2.4. Consider the decomposition (2.6) for $R = \mathbb{C}$. By [N3, Lemma 5.2], the varieties $\mathcal{M}_{\rho}/\mathbb{C}$ have no odd cohomology. By the comparison theorem for smooth proper varieties, the Poincaré polynomial $P_c(\mathcal{M}_{\rho}/\mathbb{C}, \mathbb{C})$ and the Poincaré polynomial in étale cohomology $P_c(\mathcal{M}_{\rho}/\overline{\mathbb{F}}_q, \overline{\mathbb{Q}}_l)$ coincide. In particular, the odd étale cohomology of $\mathcal{M}_{\rho}/\overline{\mathbb{F}}_q$ is zero as well. Now it is known that $\mathcal{M}(\mathbf{v}, \mathbf{w})/\overline{\mathbb{F}}_q$ is pure (see [M1, Prop. 6.2]) and strictly polynomial count (see [CBVdB, Prop. 2.2.1]). By e.g. [CBVdB, Lemma A.1] we deduce that the odd (étale) cohomology with compact support of $\mathcal{M}(\mathbf{v}, \mathbf{w})/\overline{\mathbb{F}}_q$ vanishes and that the Frobenius eigenvalues in $H^{2i}(\mathcal{M}(\mathbf{v}, \mathbf{w})/\overline{\mathbb{F}}_q, \overline{\mathbb{Q}}_l)$ are all equal to q^i . By (2.6) there is a filtration of $H^{2i}(\mathcal{M}(\mathbf{v}, \mathbf{w})/\overline{\mathbb{F}}_q, \overline{\mathbb{Q}}_l)$ whose factors are of the form $H^{2(i-u_{\rho})}(\mathcal{M}_{\rho}/\overline{\mathbb{F}}_q, \overline{\mathbb{Q}}_l)\{u_{\rho}\}$, where $\{n\}$ denotes a Tate twist and u_{ρ} is the rank of the affine fiber bundle U_{ρ} . But then all the Frobenius eigenvalues in $H^{2i}(\mathcal{M}_{\rho}/\overline{\mathbb{F}}_q, \overline{\mathbb{Q}}_l)$ are equal to q^i for all ρ and all i . Hence each $\mathcal{M}_{\rho}/\overline{\mathbb{F}}_q$ is itself polynomial count, and therefore so is $\mathcal{L}(\mathbf{v}, \mathbf{w})/\overline{\mathbb{F}}_q$. The statements concerning the purity and the counting polynomial of $\mathcal{L}(\mathbf{v}, \mathbf{w})/\overline{\mathbb{F}}_q$ follow from the decomposition of the étale cohomology of $\mathcal{L}(\mathbf{v}, \mathbf{w})$ in terms of that of the $\mathcal{M}_{\rho}$. $\square$

Corollary 2.6. *For any finite field $\mathbb{F}_q$ the following relation holds :*

$$(2.8) \quad \sum_{\mathbf{v}} q^{-d(\mathbf{v}, \mathbf{w})} |\mathcal{L}(\mathbf{v}, \mathbf{w})(\mathbb{F}_q)| z^{\mathbf{v}} = \frac{r(\mathbf{v}, q^{-1}, z)}{r(0, q^{-1}, z)}.$$

2.4. We next relate the number of points of $\Lambda_{\mathbf{v}}$ and $\mathcal{L}(\mathbf{v}, \mathbf{w})$ over finite fields. Define a stratification $\mathcal{L}(\mathbf{v}, \mathbf{w}) = \bigsqcup_{\mathbf{w}' \leq \mathbf{w}} \mathcal{L}(\mathbf{v}, \mathbf{w})_{\mathbf{w}'}$ by

$$\mathcal{L}(\mathbf{v}, \mathbf{w})_{\mathbf{w}'} = \{G_{\mathbf{v}} \cdot (\underline{x}, 0, \underline{q}) \in \mathcal{L}(\mathbf{v}, \mathbf{w}) \mid \dim(\text{Im}(\bigoplus_i q_i)) = \mathbf{w}'\}.$$

Observe that the map $(\underline{x}, 0, \underline{q}) \mapsto (\underline{x}')$ with $x'_h = q_{h''} x_h q_{h'}^{-1}$ defines an isomorphism $\mathcal{L}(\mathbf{v}, \mathbf{v})_{\mathbf{v}} \simeq \Lambda_{\mathbf{v}}$. Let $Gr_{\mathbf{w}'}^{\mathbf{w}}$ denote the Grassmannian of I -graded subspaces of W of dimension $\mathbf{w}'$. The

projection $\mathcal{L}(\mathbf{v}, \mathbf{w})_{\mathbf{w}'} \rightarrow Gr_{\mathbf{w}'}^{\mathbf{w}}$ is a fibration with fiber $\mathcal{L}(\mathbf{v}, \mathbf{w}')_{\mathbf{w}'}$. It follows that

$$(2.9) \quad |\mathcal{L}(\mathbf{v}, \mathbf{w})(\mathbb{F}_q)| = \sum_{\mathbf{w}' \leq \mathbf{w}} |Gr_{\mathbf{w}'}^{\mathbf{w}}(\mathbb{F}_q)| \cdot |\mathcal{L}(\mathbf{v}, \mathbf{w}')_{\mathbf{w}'}(\mathbb{F}_q)|.$$

Inverting (2.9) to express the number of points of $\mathcal{L}(\mathbf{v}, \mathbf{w})_{\mathbf{w}}/\mathbb{F}_q$ for all $\mathbf{w}$ in terms of the number of points of $\mathcal{L}(\mathbf{v}, \mathbf{w}')_{\mathbf{w}'}/\mathbb{F}_q$ for all $\mathbf{w}'$ yields, after a small computation, the following :

Lemma 2.7. *We have*

$$(2.10) \quad |\mathcal{L}(\mathbf{v}, \mathbf{w})_{\mathbf{w}}(\mathbb{F}_q)| = \sum_{\mathbf{w}' \leq \mathbf{w}} (-1)^{|\mathbf{w}| - |\mathbf{w}'|} q^{u(\mathbf{w}, \mathbf{w}')} |Gr_{\mathbf{w}'}^{\mathbf{w}}(\mathbb{F}_q)| \cdot |\mathcal{L}(\mathbf{v}, \mathbf{w}')_{\mathbf{w}'}(\mathbb{F}_q)|,$$

where $u(\mathbf{w}, \mathbf{w}') = \sum_i (w_i - w'_i)(w_i - w'_i - 1)/2$.

Corollary 2.8.

$$(2.11) \quad |\Lambda_{\mathbf{v}}(\mathbb{F}_q)| = \sum_{\mathbf{w}' \leq \mathbf{v}} (-1)^{|\mathbf{v}| - |\mathbf{w}'|} q^{u(\mathbf{v}, \mathbf{w}')} |Gr_{\mathbf{w}'}^{\mathbf{v}}(\mathbb{F}_q)| \cdot |\mathcal{L}(\mathbf{v}, \mathbf{w}')_{\mathbf{w}'}(\mathbb{F}_q)|.$$

2.5. We may now proceed to the proof of Theorem 1.1. It is essentially a direct computation using (2.11) together with (2.8). For this, we consider the series

$$(2.12) \quad T_{\mathbf{w}}(z) = \sum_{\mathbf{v}} \frac{|\mathcal{L}(\mathbf{v}, \mathbf{w})_{\mathbf{w}}(\mathbb{F}_q)|}{|G_{\mathbf{w}}(\mathbb{F}_q)|} q^{\langle \mathbf{v}, \mathbf{v} \rangle} z^{\mathbf{v}} \in \frac{|\Lambda_{\mathbf{w}}(\mathbb{F}_q)|}{|G_{\mathbf{w}}(\mathbb{F}_q)|} q^{\langle \mathbf{w}, \mathbf{w} \rangle} z^{\mathbf{w}} + \bigoplus_{\mathbf{v} > \mathbf{w}} \mathbb{C} z^{\mathbf{v}}.$$

Using Lemma 2.7 and Corollary 2.6 we have

$$\begin{aligned} T_{\mathbf{w}}(z) &= \sum_{\mathbf{w}' \leq \mathbf{w}} \left\{ (-1)^{|\mathbf{w}| - |\mathbf{w}'|} \frac{|Gr_{\mathbf{w}'}^{\mathbf{w}}(\mathbb{F}_q)|}{|G_{\mathbf{w}}(\mathbb{F}_q)|} q^{u(\mathbf{w}, \mathbf{w}')} \sum_{\mathbf{v}} |\mathcal{L}(\mathbf{v}, \mathbf{w}')_{\mathbf{w}'}(\mathbb{F}_q)| q^{\langle \mathbf{v}, \mathbf{v} \rangle} z^{\mathbf{v}} \right\} \\ &= \sum_{\mathbf{w}' \leq \mathbf{w}} (-1)^{|\mathbf{w}| - |\mathbf{w}'|} \frac{|Gr_{\mathbf{w}'}^{\mathbf{w}}(\mathbb{F}_q)|}{|G_{\mathbf{w}}(\mathbb{F}_q)|} q^{u(\mathbf{w}, \mathbf{w}')} \frac{r(\mathbf{w}', q^{-1}, q^{\mathbf{w}'} z)}{r(0, q^{-1}, q^{\mathbf{w}'} z)}, \end{aligned}$$

where by convention $(q^{\mathbf{x}} z)^{\mathbf{y}} = q^{\mathbf{x} \cdot \mathbf{y}} z^{\mathbf{y}}$. Expanding in powers of z , we have

$$\begin{aligned} \frac{r(\mathbf{w}', q^{-1}, q^{\mathbf{w}'} z)}{r(0, q^{-1}, q^{\mathbf{w}'} z)} &= \\ &= \left\{ \sum_{\boldsymbol{\tau}} q^{\mathbf{w}' \cdot |\boldsymbol{\tau}|_{>1}} |z|^{\boldsymbol{\tau}} X(\boldsymbol{\tau}, q) \right\} \cdot \left\{ \sum_{l \geq 0} (-1)^l \sum_{\substack{\boldsymbol{\tau}^{(1)}, \dots, \boldsymbol{\tau}^{(l)} \\ \boldsymbol{\tau}^{(j)} \neq 0}} q^{\mathbf{w}' \cdot (\sum_j |\boldsymbol{\tau}^{(j)}|)} z^{\sum_j |\boldsymbol{\tau}^{(j)}|} \prod_j X(\boldsymbol{\tau}^{(j)}, q) \right\} \end{aligned}$$

It follows that

$$(2.13) \quad T_{\mathbf{w}}(z) = \frac{1}{|G_{\mathbf{w}}(\mathbb{F}_q)|} \sum_{l \geq 0} (-1)^l \sum_{\substack{\boldsymbol{\tau}^{(1)}, \dots, \boldsymbol{\tau}^{(l)} \\ \boldsymbol{\tau}^{(j)} \neq 0}} K_{\mathbf{w}}^{(l)}(\boldsymbol{\tau}^{(1)}, \dots, \boldsymbol{\tau}^{(l)}) \cdot z^{\sum |\boldsymbol{\tau}^{(j)}|} \prod_j X(\boldsymbol{\tau}^{(j)}, q)$$

where

$$K_{\mathbf{w}}^{(l)}(\boldsymbol{\tau}^{(1)}, \dots, \boldsymbol{\tau}^{(l)}) = \sum_{\mathbf{w}' \leq \mathbf{w}} (-1)^{|\mathbf{w}| - |\mathbf{w}'|} q^{u(\mathbf{w}, \mathbf{w}')} |Gr_{\mathbf{w}'}^{\mathbf{w}}(\mathbb{F}_q)| (q^{|\mathbf{w}'| \cdot |\boldsymbol{\tau}_{>1}^{(1)}|} - q^{|\mathbf{w}'| \cdot |\boldsymbol{\tau}^{(1)}|}) q^{\mathbf{w}' \cdot (\sum_{j>1} |\boldsymbol{\tau}^{(j)}|)}$$

Replacing $\mathbf{w}'$ by $\mathbf{w} - \mathbf{w}'$ and setting $\mathbf{1} = (1, \dots, 1) \in \mathbb{N}^I$ we may rewrite $K_{\mathbf{w}}^{(l)}(\tau^{(1)}, \dots, \tau^{(l)})$ as

$$\begin{aligned} K_{\mathbf{w}}^{(l)}(\tau^{(1)}, \dots, \tau^{(l)}) &= \\ &= q^{|\mathbf{w}| \cdot (|\tau_{>1}^{(1)}| + \sum_{j>1} |\tau^{(j)}|)} \left(\sum_{\mathbf{w}' \leq \mathbf{w}} (-1)^{|\mathbf{w}'|} q^{\frac{1}{2} \mathbf{w}' \cdot (\mathbf{w}' - \mathbf{1})} |Gr_{\mathbf{w}'}^{\mathbf{w}}(\mathbb{F}_q)| q^{-|\mathbf{w}'| \cdot (|\tau_{>1}^{(1)}| + \sum_{j>1} |\tau^{(j)}|)} \right) \\ &\quad - q^{|\mathbf{w}| \cdot (\sum_j |\tau^{(j)}|)} \left(\sum_{\mathbf{w}' \leq \mathbf{w}} (-1)^{|\mathbf{w}'|} q^{\frac{1}{2} \mathbf{w}' \cdot (\mathbf{w}' - \mathbf{1})} |Gr_{\mathbf{w}'}^{\mathbf{w}}(\mathbb{F}_q)| q^{-|\mathbf{w}'| \cdot (\sum_j |\tau^{(j)}|)} \right). \end{aligned}$$

Now we use the following identity, whose proof is left to the reader. For any $w \in \mathbb{N}$ we have

$$\sum_{j=0}^w (-1)^{w'} q^{w'(w'-1)/2 - aw'} |Gr_{w'}^w(\mathbb{F}_q)| = \begin{cases} 0 & \text{if } a = 0, 1, \dots, d-1 \\ (1-q)^{-1} \dots (1-q^{-w})^{-1} & \text{if } a = d \end{cases}.$$

This implies that $K_{\mathbf{w}}^{(l)}(\tau^{(1)}, \dots, \tau^{(l)}) = 0$ unless $\sum_i |\tau^{(i)}| - \mathbf{w} \in \mathbb{N}^I$, and that for $\sum_i |\tau^{(i)}| = \mathbf{w}$ we have

$$K_{\mathbf{w}}^{(l)}(\tau^{(1)}, \dots, \tau^{(l)}) = -|G_{\mathbf{w}}(\mathbb{F}_q)|.$$

Comparing the coefficients of $z^{\mathbf{w}}$ in (2.13) and (2.12) we obtain the equality

$$\frac{|\Lambda_{\mathbf{w}}(\mathbb{F}_q)|}{|G_{\mathbf{w}}(\mathbb{F}_q)|} q^{\langle \mathbf{w}, \mathbf{w} \rangle} = \sum_l (-1)^{l+1} \left\{ \sum_{\tau^{(1)}, \dots, \tau^{(l)}} \prod_j X(\tau^{(j)}, q) \right\}$$

where the sums runs over all tuples $(\tau^{(1)}, \dots, \tau^{(l)})$ of nonzero partitions satisfying $\sum_j |\tau^{(j)}| = \mathbf{w}$. Summing over all $\mathbf{w}$ we finally obtain

$$\sum_{\mathbf{w}} \frac{|\Lambda_{\mathbf{w}}(\mathbb{F}_q)|}{|G_{\mathbf{w}}(\mathbb{F}_q)|} q^{\langle \mathbf{w}, \mathbf{w} \rangle} z^{\mathbf{w}} = \frac{1}{1 + \sum_{\tau \neq 0} X(\tau, q) z^{|\tau|}} = \frac{1}{r(0, q^{-1}, z)}.$$

The theorem is now a consequence of Hua's formula

$$r(0, q^{-1}, z) = \text{Exp} \left(\frac{1}{q^{-1} - 1} \sum_{\mathbf{d}} A_{\mathbf{d}}(q^{-1}) z^{\mathbf{d}} \right) = \text{Exp} \left(\frac{1}{q - 1} \sum_{\mathbf{d}} \bar{A}_{\mathbf{d}}(q) q^{\langle \mathbf{d}, \mathbf{d} \rangle} z^{\mathbf{d}} \right)^{-1}.$$

□

3. FACTORIZATION OF $\lambda_Q(q, z)$ AND THE STRATA IN $\Lambda_{\mathbf{d}}$.

3.1. Let Q_J be the full subquiver of Q corresponding to a subset of vertices $J \subset I$. All the varieties associated to Q_J instead of Q will be denoted with a superscript J . If $\mathbf{d} \in \mathbb{N}^J$ then of course $\Lambda_{\mathbf{d}}^J \simeq \Lambda_{\mathbf{d}}$ so that there is a factorization

$$(3.1) \quad \lambda_Q(q, z) = \lambda_{Q_J}(q, z) \cdot \lambda_{Q \setminus Q_J}(q, z)$$

where by definition

$$\lambda_{Q \setminus Q_J}(q, z) = \text{Exp} \left(\frac{1}{q - 1} \sum_{\substack{\mathbf{d} \\ \text{supp}(\mathbf{d}) \not\subseteq J}} \bar{A}_{\mathbf{d}}(q) q^{\langle \mathbf{d}, \mathbf{d} \rangle} z^{\mathbf{d}} \right).$$

The Fourier modes of $\lambda_{Q \setminus Q_J}(q, z)$ count the (orbifold) number of points of some natural subvarieties in the $\Lambda_{\mathbf{d}}$ first considered by Lusztig. These subvarieties are defined as follows. Let $K = \{k \in H \sqcup H^* \mid k' \in I \setminus J, k'' \in J\}$. For a pair $(\mathbf{d}, \mathbf{n}) \in \mathbb{N}^I \times \mathbb{N}^J$ such that $\mathbf{d} - \mathbf{n} \in \mathbb{N}^I$, we set

$$\Lambda_{\mathbf{d}, \mathbf{n}} = \left\{ (x_h, x_{h^*})_h \in \Lambda_{\mathbf{d}} \mid \text{codim} \left(\bigoplus_{k \in K} k \right) = \mathbf{n} \right\}.$$

Each $\Lambda_{\mathbf{d};\mathbf{n}}$ is a locally closed subvariety in $\Lambda_{\mathbf{d}}$ and we have a stratification

$$\Lambda_{\mathbf{d}} = \bigsqcup_{\mathbf{n}} \Lambda_{\mathbf{d};\mathbf{n}}.$$

There is natural map of stacks

$$p_{\mathbf{d},\mathbf{n}} : [\Lambda_{\mathbf{d};\mathbf{n}}/G_{\mathbf{d}}] \rightarrow [\Lambda_{\mathbf{d};0}/G_{\mathbf{d}}] \times [\Lambda_{\mathbf{n}}^J/G_{\mathbf{n}}]$$

given by assigning to a representation $M = (x_h, x_{h^*})_h$ in $\Lambda_{\mathbf{d};\mathbf{n}}$ the pair $(F, M/F)$ where F is the subrepresentation of M (for the doubled quiver $\overline{Q}$) generated by $\bigoplus_{i \notin J} V_i$. The following is proved in [L1, Sect.12] and plays a key role in the geometric construction of the crystal graph. The proof in *loc. cit* is given in the case that J is reduced to a single vertex, but the proof is the same in general, see e.g. [B].

Proposition 3.1. *The map $p_{\mathbf{d},\mathbf{n}}$ is a (not necessarily representable) affine fibration of dimension $(\mathbf{d} - \mathbf{n}, \mathbf{n})$.*

Set

$$\lambda_{Q;0}(q, z) = \sum_{\mathbf{d}} \frac{|\Lambda_{\mathbf{d};0}(\mathbb{F}_q)|}{|G_{\mathbf{d}}(\mathbb{F}_q)|} q^{\langle \mathbf{d}, \mathbf{d} \rangle} z^{\mathbf{d}}.$$

Then from Proposition 3.1 we easily deduce the identity

$$(3.2) \quad \lambda_Q(q, z) = \lambda_{Q_J}(q, z) \cdot \lambda_{Q;0}(q, z).$$

Corollary 3.2. *The following equality holds*

$$\lambda_{Q;0}(q, z) = \lambda_{Q \setminus Q_J}(q, z).$$

Acknowledgements

I would like to thank A. Chambert-Loir, T. Hausel, E. Letellier, H. Nakajima and Y. Soibelman for useful discussions and correspondence. As explained to us by Y. Soibelman, a formula similar to (1.2) has been obtained independently in some recent joint work of his and Kontsevich.

REFERENCES

- [B] T. Bozec, in preparation.
- [CBVdB] W. Crawley-Boevey, M. Van den Bergh, *Absolutely indecomposable representations and Kac-Moody Lie algebras. With an appendix by Hiraku Nakajima*, Invent. Math. **155** (2004), no. 3, 537–559.
- [FT] W. Feit, N. J. Fine, *Pairs of commuting matrices over a finite field*, Duke Math. J. **27** 1960 91–94.
- [Ha] T. Hausel, *Kac’s conjecture from Nakajima quiver varieties*, Invent. Math. **181** (2010), no. 1, 21–37.
- [HLR-V] T. Hausel, E. Letellier, F. Villegas-Rodriguez, *Positivity of Kac polynomials and DT-invariants for quivers*, arXiv:1204.2375 (2012).
- [He] W. Hesselink, *Concentration under actions of algebraic groups*, Séminaire d’Algèbre Paul Dubreil et Marie-Paule Malliavin, Lecture Notes in Mathematics No. **867**, 1981, pp 55–89.
- [Hu] J. Hua, *Counting representations of quivers over finite fields*, J. Algebra **226**, 1011–1033 (2000)
- [K] V. Kac, *Root systems, representations of quivers and invariant theory*, In: Invariant Theory, Montecatini, 1982. Lecture Notes in Mathematics, vol. **996**, pp. 74–108. Springer, Berlin (1983).
- [KS] M. Kashiwara, T. Saito, *Geometric construction of crystal bases*, Duke Math. J. **89** (1997), no. 1, 9–36.
- [L1] G. Lusztig, *Quivers, perverse sheaves, and quantized enveloping algebras*, J. Amer. Math. Soc. **4** (1991), no. 2, 365–421.
- [L2] G. Lusztig, *Canonical bases arising from quantized enveloping algebras. II*, Common trends in mathematics and quantum field theories (Kyoto, 1990). Progr. Theoret. Phys. Suppl. No. **102** (1990), 175–201 (1991).
- [M1] S. Mozgovoy, *Fermionic forms and quiver varieties*, arXiv:math/0610084 (2006).
- [M2] S. Mozgovoy, *Motivic Donaldson-Thomas invariants and McKay correspondence*, arXiv:1107.6044 (2011).
- [N1] H. Nakajima, *Quiver varieties and Kac-Moody algebras*, Duke Math. J. **91**(3), 515–560 (1998).

- [N2] H. Nakajima, *Quiver varieties and finite dimensional representations of quantum affine algebras*, J. Amer. Math. Soc. **14** (2001), no. 1, 145–238.
- [N3] H. Nakajima, *Quiver varieties and t -analogs of q -characters of quantum affine algebras*, Ann. of Math. (2) **160** (2004), no. 3, 1057–1097.
- [R-V] F. Villegas-Rodriguez, *Counting colorings on varieties*, Publ. Mat. 2007, Proceedings of the Primeras Jornadas de Teoría de Números, 209–220.
- [S] O. Schiffmann, *Lectures on canonical and crystal bases of Hall algebras*, arXiv:0910.4460 (2009), to appear in Sémin. et Cong. (SMF).

O. Schiffmann, `olivier.schiffmann@math.u-psud.fr`,
Département de Mathématiques, Université de Paris-Sud, Bâtiment 425 91405 Orsay Cedex, FRANCE.